

VPS35 Antibody

VPS35 Antibody, Clone 11H10 Catalog # ASM10695

Specification

VPS35 Antibody - Product Information

Primary Accession
Other Accession
Host
Clonality
Target/Specificity

Q96QK1 NP_060676.2 Mouse Monoclonal

Blue Ice or 4ºC

Other Names

VPS35

Vacuolar protein sorting-associated protein 35, MEM3, PARK17, VPS35 retromer complex component, maternal-embryonic 3, vesicle protein sortin 35, TCCCTA00141, FLJ10752

Immunogen

Full length recombinant human VSP35

Purification

Protein G Purified

Storage -20°C

Storage Buffer

PBS pH 7.4, 50% glycerol, 0.09% Sodium azide *Storage buffer may change when conjugated

Shipping Temperature

Certificate of Analysis

A 1:1000 dilution of SMC-606 was sufficient for detection of VPS35 in 10 μ g of SH-SY5Y by ECL immunoblot analysis using Goat Anti-Mouse IgG:HRP as the secondary antibody.

Cellular Localization

Endosome | Lysosome | Vesicles | Cytoplasm | Membrane


VPS35 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

VPS35 Antibody - Images






Immunohistochemistry analysis using Mouse Anti-VPS35 Monoclonal Antibody, Clone 11H10 (ASM10695). Tissue: Intestinal Cancer. Species: Human. Primary Antibody: Mouse Anti-VPS35 Monoclonal Antibody (ASM10695) at 1:100 for Overnight at 4C, then 30 min at 37C. Secondary Antibody: Goat Anti-Mouse IgG (H+L): FITC for 45 min at 37C. Counterstain: DAPI for 3 min at RT. Magnification: 20X.




Immunoprecipitation analysis using Mouse Anti-VPS35 Monoclonal Antibody, Clone 11H10 (ASM10695). Tissue: A549 cells. Species: Human. Primary Antibody: Mouse Anti-VPS35 Monoclonal Antibody (ASM10695). 500 μ L cell culture supernatants were incubated with 10 μ L of Protein A/G resin beads for 1 hour at 4°C. ASM10695 clone 11H10 depletes VPS35 from the A549 cell extract.






Immunoprecipitation analysis using Mouse Anti-VPS35 Monoclonal Antibody, Clone 11H10 (ASM10695). Tissue: A549 cells. Species: Human. Primary Antibody: Mouse Anti-VPS35 Monoclonal Antibody (ASM10695).




Western Blot analysis of Human, Mouse A549, MEF showing detection of VPS35 protein using Mouse Anti-VPS35 Monoclonal Antibody, Clone 11H10 (ASM10695). Lane 1: Molecular Weight Ladder. Lane 2: VPS35 KO A549 cells. Lane 3: mouse embryonic fibroblast cells.. Load: 8 µg each A549 and MEF. Primary Antibody: Mouse Anti-VPS35 Monoclonal Antibody (ASM10695) at 1:5 (tissue culture supernatant). Secondary Antibody: Donkey anti-mouse IRDye 800CW at 1:25000 in TBS-T.






Western Blot analysis of Human SH-SY5Y showing detection of VPS35 protein using Mouse Anti-VPS35 Monoclonal Antibody, Clone 11H10 (ASM10695). Lane 1: Molecular Weight Ladder. Lane 2: SH-SY5Y (10 ug). Load: 10 μ g. Block: 5% Skim Milk powder in TBST. Primary Antibody: Mouse Anti-VPS35 Monoclonal Antibody (ASM10695) at 1:1000 for 2 hours at RT with shaking. Secondary Antibody: Goat anti-mouse IgG:HRP at 1:4000 for 1 hour at RT with shaking. Color Development: Chemiluminescent for HRP (Moss) for 5 min in RT.



Immunoprecipitation analysis using Mouse Anti-VPS35 Monoclonal Antibody, Clone 11H10 (ASM10695). Tissue: A549 cells. Species: Human. Primary Antibody: Mouse Anti-VPS35 Monoclonal Antibody (ASM10695). Three amounts of ASM10695 (3, 1 and 0.3 ug) were non-covalently coupled to 10uL of A/G sepharose beads for 1 hour at 4°C and next incubated with 250ug of A549 lysate for 2 hours at 4°C.





Immunoprecipitation analysis using Mouse Anti-VPS35 Monoclonal Antibody, Clone 11H10 (ASM10695). Tissue: embryonic fibroblast. Species: Mouse. Primary Antibody: Mouse Anti-VPS35 Monoclonal Antibody (ASM10695). Three amounts of ASM10695 (3, 1 and 0.3 ug) were non-covalently coupled to 10uL of A/G sepharose beads for 1 hour at 4°C and next incubated with 250ug of MEF lysate for 2 hours at 4°C.

VPS35 Antibody - Background

Vacuolar Protein Sorter-35 (VPS35) is a component of the retromer complex, which is essential for endosome-to-Golgi retrieval of membrane proteins. VPS35 mutations such as D620N have been linked to Parkinson's Disease (PD) (1,2) and affect retromer function, protein homeostasis, and mitochondria (3).

VPS35 Antibody - References

- 1. Vilarino-Guell, C. et al. (2011) Am J Hum Genet 89:162-167
- 2. Zimprich, A. et al. (2011) Am J Hum Genet 89:168-175
- 3. Rahman, A.A., Morrison, B.E. (2019) Neurosci 401:1-10.