

Mouse IgG3 isotype control Rhodamine
Monoclonal MG3 IgG3 , Rhodamine (TRITC)
Catalog # ASR1108

Specification

Mouse IgG3 isotype control Rhodamine - Product Information

Description	MOUSE IgG3 isotype control Rhodamine conjugated Rhodamine (TRITC)
Conjugate	2.6 moles Rhodamine (TRITC) per mole of Mouse IgG3
FP Value	Monoclonal
Clonality	FC
Application	FlowCytometry 1:1000-1:5000
Application Note	Lyophilized
Physical State	IgG3
Host Isotype	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Buffer	Mouse
Species of Origin	100 µL
Reconstitution Volume	Restore with deionized water (or equivalent)
Reconstitution Buffer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Stabilizer	0.01% (w/v) Sodium Azide
Preservative	

Mouse IgG3 isotype control Rhodamine - Additional Information

Shipping Condition

Ambient

Purity

This product was prepared from normal serum by delipidation, salt fractionation, ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunolectrophoresis resulted in a single precipitin arc against anti-Mouse IgG3 and anti-Mouse Serum.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Mouse IgG3 isotype control Rhodamine - Protein Information

Mouse IgG3 isotype control Rhodamine - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Mouse IgG3 isotype control Rhodamine - Images

Mouse IgG3 isotype control Rhodamine - Background

Isotype controls are important for Flow Cytometry and have no specificity for target cells within a particular experiment. Their purpose is to confirm the specificity of primary antibody binding that it is not a result of non-specific Fc receptor binding to cells or other cellular protein interactions. Isotype controls need to be matched to the specific primary Abs (species and isotype, including heavy and light chains) being used.