

Anti-Donkey IgG (H&L) Secondary Antibody

Goat Polyclonal, Unconjugated Catalog # ASR1375

Specification

Physical State

Anti-Donkey IgG (H&L) Secondary Antibody - Product Information

Description Anti-DONKEY IgG (H&L) (GOAT) Antibody

Host Goat

Conjugate
Target Species
Clonality
Application

Unconjugated
Donkey
Polyclonal
,1,10,15,

Application Note ELISA 1:20,000-1:100,000;Western Blot

1:2,000-1:10,000;Immunochemistry

1:1,000-1:5,000 Lyophilized Antiserum

Host Isotype Antiserum
Target Isotype IgG (H&L)

Buffer 0.01 M Sodium Phosphate, 0.15 M Sodium

Chloride, pH 7.2

Immunogen Donkey IgG whole molecule

Reconstitution Volume 2.0 m

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer None

Preservative 0.01% (w/v) Sodium Azide

Anti-Donkey IgG (H&L) Secondary Antibody - Additional Information

Shipping Condition

Ambient

Purity

This product was prepared from monospecific antiserum by a delipidation and defibrination. Assay by immunoelectrophoresis resulted in a single precipitin arc against Donkey IgG and Donkey Serum.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Donkey IgG (H&L) Secondary Antibody - Protein Information



Anti-Donkey IgG (H&L) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-Donkey IgG (H&L) Secondary Antibody - Images