

Swine IgG F(ab')2 Peroxidase

Catalog # ASR1513

Specification

Conjugate

Buffer

Stabilizer

Preservative

Physical State

Species of Origin

Reconstitution Volume

Reconstitution Buffer

Host Isotype

Swine IgG F(ab')2 Peroxidase - Product Information

Description SWINE IgG F(ab')2 fragment

Peroxidase conjugated Peroxidase (Horseradish)

Lyophilized IgG F(ab')2

0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Swine 1.0 mL

Restore with deionized water (or

equivalent)

10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

0.01% (w/v) Gentamicin Sulfate. Do NOT

add Sodium Azide!

Swine IgG F(ab')2 Peroxidase - Additional Information

Shipping Condition

Ambient

Purity

This product was prepared from normal serum by delipidation, salt fractionation, ion exchange chromatography followed by pepsin digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Swine IgG, anti-Swine IgG F(ab')2 and anti-Swine Serum. No reaction was observed against anti-Swine IgG F(c) or anti-Pepsin.

Storage Condition


Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Swine IgG F(ab')2 Peroxidase - Protein Information

Swine IgG F(ab')2 Peroxidase - Protocols





Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Swine IgG F(ab')2 Peroxidase - Images