

**Anti-Cat IgG (H&L) (Rhodamine Conjugated) Secondary Antibody**  
**Goat Polyclonal, Rhodamine (TRITC)**  
**Catalog # ASR1548**

**Specification**

**Anti-Cat IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Product Information**

|                       |  |
|-----------------------|--|
| Description           | Anti-CAT IgG (H&L) (GOAT) Antibody<br>Rhodamine Conjugated                                     |
| Host                  | Goat   |
| Conjugate             | Rhodamine (TRITC)  |
| FP Value              | 3.0 moles Rhodamine (TRITC) per mole of IgG  |
| Target Species        | Cat  |
| Clonality             | Polyclonal   |
| Application           | IF, FC   |
| Application Note      | FLISA 1:10,000-1:50,000;IF Microscopy<br>1:1,000-1:5,000;FlowCytometry<br>1:500-1:2,500        |
| Physical State        | Lyophilized  |
| Host Isotype          | IgG  |
| Target Isotype        | IgG (H&L)  |
| Buffer                | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2<br>Cat IgG whole molecule<br>1.0 mL |
| Immunogen             | Restore with deionized water (or equivalent)   |
| Reconstitution Volume | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free                         |
| Reconstitution Buffer | 0.01% (w/v) Sodium Azide   |
| Stabilizer            |  |
| Preservative          |  |

**Anti-Cat IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Additional Information**

**Shipping Condition**

Ambient

**Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Cat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Cat IgG and Cat Serum.

**Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

**Anti-Cat IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Protein Information****Anti-Cat IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-Cat IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Images****Anti-Cat IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Background**

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.