

Anti-Goat IgG (H&L) (Biotin Conjugated) Secondary Antibody
Donkey Polyclonal, Biotin
Catalog # ASR1568**Specification****Anti-Goat IgG (H&L) (Biotin Conjugated) Secondary Antibody - Product Information**

Description	Anti-GOAT IgG (H&L) (DONKEY) Antibody
Host	Biotin Conjugated
Conjugate	Donkey
FP Value	Biotin
Target Species	10-20 moles Biotin per mole of IgG
Reactivity	Goat
Clonality	Goat
Application	Polyclonal
Application Note	,1,10,15, ELISA 1:140,000;Western Blot 1:2,000-1:10,000;Immunochemistry 1:1,000-1:5,000
Physical State	Lyophilized
Host Isotype	IgG
Target Isotype	IgG (H&L)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Anti-Goat IgG (h&L) was produced by repeated immunization with goat IgG whole molecule in donkey.
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Sodium Azide

Anti-Goat IgG (H&L) (Biotin Conjugated) Secondary Antibody - Additional Information**Shipping Condition**

Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Donkey Serum, Goat IgG and Goat Serum.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Goat IgG (H&L) (Biotin Conjugated) Secondary Antibody - Protein Information**Anti-Goat IgG (H&L) (Biotin Conjugated) Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Goat IgG (H&L) (Biotin Conjugated) Secondary Antibody - Images**Anti-Goat IgG (H&L) (Biotin Conjugated) Secondary Antibody - Background**

Anti-Goat IgG (H&L) antibody generated in donkey detects specifically goat IgG. This secondary antibody anti-Goat is ideal for investigators who routinely perform titration assays, western-blot, immunoprecipitation and more generally immunoassays.