

**Fab Anti-Golden Syrian Hamster IgG (H&L) (Rhodamine Conjugated) Secondary Antibody**  
**Goat Polyclonal, Rhodamine (TRITC)**  
**Catalog # ASR1711****Specification****Fab Anti-Golden Syrian Hamster IgG (H&L) (Rhodamine Conjugated) Secondary Antibody**  
**- Product Information**

Description	<b>F(ab) Anti-GS HAMSTER IgG (H&amp;L) (GOAT) Antibody Rhodamine Conjugated</b>
Host	<b>Goat</b>
Conjugate	<b>Rhodamine (TRITC)</b>
FP Value	<b>2.0 moles Rhodamine (TRITC) per mole of IgG F(ab)</b>
Target Species	<b>Golden Syrian Hamster</b>
Clonality	<b>Polyclonal</b>
Application	<b>IF, FC</b>
Application Note	<b>FLISA 1:10,000-1:50,000;IF Microscopy 1:1,000-1:5,000;FlowCytometry 1:500-1:2,500</b>
Physical State	<b>Lyophilized</b>
Host Isotype	<b>IgG F(ab)</b>
Target Isotype	<b>IgG (H&amp;L)</b>
Buffer	<b>0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2</b>
Immunogen	<b>Golden Syrian Hamster IgG whole molecule</b>
Reconstitution Volume	<b>1.0 mL</b>
Reconstitution Buffer	<b>Restore with deionized water (or equivalent)</b>
Stabilizer	<b>10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free</b>
Preservative	<b>0.01% (w/v) Sodium Azide</b>

**Fab Anti-Golden Syrian Hamster IgG (H&L) (Rhodamine Conjugated) Secondary Antibody**  
**- Additional Information****Shipping Condition**

Ambient

**Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Golden Syrian Hamster IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein and anti-Goat Serum. No reaction was observed against anti-Papain or anti-Goat IgG F(c).

**Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted

liquid. Dilute only prior to immediate use.

**Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

**Fab Anti-Golden Syrian Hamster IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Protein Information****Fab Anti-Golden Syrian Hamster IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Fab Anti-Golden Syrian Hamster IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Images****Fab Anti-Golden Syrian Hamster IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Background**

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.