

Anti-GOAT IgG (H&L) (Peroxidase Conjugated) Secondary Antibody
Rabbit Polyclonal, Peroxidase (Horseradish)
Catalog # ASR1770**Specification****Anti-GOAT IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Product Information**

Description	Anti-GOAT IgG (H&L) (RABBIT) Antibody Peroxidase Conjugated
Host	Rabbit
Conjugate	Peroxidase (Horseradish)
Target Species	Goat
Clonality	Polyclonal
Application	,1,10,15,
Application Note	ELISA 1:300,000;Western Blot 1:5,000-1:20,000;Immunochemistry 1:1,000-1:5,000
Physical State	Lyophilized
Host Isotype	IgG
Target Isotype	IgG (H&L)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Goat IgG whole molecule
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Anti-GOAT IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Additional Information**Shipping Condition**

Ambient

Purity

Secondary Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Goat IgG and Goat Serum.

Storage Condition

Store Secondary Antibody at 4° C prior to restoration. For extended storage aliquot antibody and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.


Anti-GOAT IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Protein Information

Anti-GOAT IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-GOAT IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Images



Western Blot of Rabbit anti-GOAT IgG (H&L) Antibody Peroxidase Conjugated Lane 1: Goat IgG Load: 100 ng per lane Secondary antibody: GOAT IgG (H&L) Antibody Peroxidase Conjugated at 1:1,000 for 60 min at RT Block: MB-070 for 30 min at RT

Anti-GOAT IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Background

HRP Conjugated Secondary Antibodies are ideal for Western Blotting, Immunohistochemistry, ELISA as well as other antibody detection methods.