

Anti-Horse IgG F(c) Secondary Antibody

Rabbit Polyclonal, Unconjugated Catalog # ASR1774

Specification

Anti-Horse IgG F(c) Secondary Antibody - Product Information

Description Anti-HORSE IgG F(c) (RABBIT) Antibody

Host Rabbit

Conjugate Unconjugated

Target Species
Clonality
Application
Horse
Polyclonal
,1,10,15,

Application Note ELISA 1:2,000,000; Western Blot

1:2,000-1:10,000;Immunochemistry

1:1,000-1:5,000

Physical State Liquid (sterile filtered)

Host Isotype IgG
Target Isotype IgG F(c)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Immunogen Horse IgG F(c) fragment

Stabilizer None

Preservative 0.01% (w/v) Sodium Azide

Anti-Horse IgG F(c) Secondary Antibody - Additional Information

Shipping Condition

Wet Ice

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Horse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Horse IgG, Horse IgG F(c) and Horse Serum. No reaction was observed against Horse IgG F(ab')2.

Storage Condition

Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Horse IgG F(c) Secondary Antibody - Protein Information



Tel: 858.875.1900 Fax: 858.875.1999



Anti-Horse IgG F(c) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-Horse IgG F(c) Secondary Antibody - Images