

# Anti-Rat IgG (H&L) (Peroxidase Conjugated) Secondary Antibody

Sheep Polyclonal, Peroxidase (Horseradish)
Catalog # ASR1833

#### **Specification**

# Anti-Rat IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Product Information

Description Anti-RAT IgG (H&L) (SHEEP) Antibody

**Peroxidase Conjugated** 

Host Sheep

Conjugate Peroxidase (Horseradish)

Target Species Rat

Clonality Polyclonal Application WB, E, IC

Application Note ELISA 1:10,000-1:50,000; Western Blot

1:1,000-1:10,000;Immunochemistry

1:500-1:2,500 Lyophilized

Physical State
Host Isotype
IgG

Target Isotype IgG (H&L)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2 Rat IgG whole molecule

Immunogen Rat IgG

Reconstitution Volume 1.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Gentamicin Sulfate. Do NOT

add Sodium Azide!

# Anti-Rat IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Additional Information

#### **Shipping Condition**

**Ambient** 

### **Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Sheep Serum, Rat IgG and Rat Serum.

#### **Storage Condition**

Store vial at  $4^{\circ}$  C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at  $4^{\circ}$  C as an undiluted liquid. Dilute only prior to immediate use.

#### **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.



### Anti-Rat IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Protein Information

## Anti-Rat IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

### Anti-Rat IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Images



Western Blot of Anti-Rat IgG (H&L) (SHEEP) Antibody . Lane M: 3  $\mu$ l Molecular Ladder. Lane 1: Rat IgG whole molecule . Lane 2: Rat IgG F(c) Fragment . Lane 3: Rat IgG F(ab) Fragment . Lane 4: Rat IgM Whole Molecule . Lane 5: Rat Serum . All samples were reduced. Load: 50 ng per Iane. Block: MB-070 for 30 min at RT. Primary Antibody: Anti-Rat IgG (H&L) (SHEEP) Antibody 1:1,000 for 60 min at RT. Secondary Antibody: Anti-Sheep IgG (DONKEY) Peroxidase Conjugated Antibody 1:40,000 in MB-070 for 30 min at RT. Predicted/Obsevered Size: 25 and 55 kDa for Rat IgG and Serum, 25 kDa for F(c) and F(ab), 78 and 25 kDa for IgM. Rat F(c) migrates slightly higher.