

F(ab')2 Anti-Bovine IgG (H&L) (Biotin Conjugated) Secondary Antibody

Rabbit Polyclonal, Biotin Catalog # ASR1842

Specification

F(ab')2 Anti-Bovine IgG (H&L) (Biotin Conjugated) Secondary Antibody - Product Information

Description F(ab')2 Anti-BOVINE IgG [H&L]

Host Rabbit Conjugated Biotin Conjugated Biotin

Target Species
Clonality
Application
Biotin
Bovine
Polyclonal
1,2,10,

Application Note ELISA 1:10,000-1:50,000; Western Blot

1:1,000-1:5,000;Immunohistochemistry

Physical State
Host Isotype
Target Isotype

1:500-1:2,000
Lyophilized
IgG F(ab')2
IgG (H&L)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2
Bovine IgG whole molecule

Reconstitution Volume 500 μL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/ml Polyethylene Glycol (PEG-8000)

Preservative 0.01% (w/v) Sodium Azide

F(ab')2 Anti-Bovine IgG (H&L) (Biotin Conjugated) Secondary Antibody - Additional Information

Shipping Condition

Ambient

Immunogen

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Bovine IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Rabbit Serum, Bovine IgG and Bovine Serum. No reaction was observed against anti-Pepsin or anti-Rabbit IgG F(c).

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note



Tel: 858.875.1900 Fax: 858.875.1999

This product is for research use only and is not intended for therapeutic or diagnostic applications.

F(ab')2 Anti-Bovine IgG (H&L) (Biotin Conjugated) Secondary Antibody - Protein Information

F(ab')2 Anti-Bovine IgG (H&L) (Biotin Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

F(ab')2 Anti-Bovine IgG (H&L) (Biotin Conjugated) Secondary Antibody - Images

F(ab')2 Anti-Bovine IgG (H&L) (Biotin Conjugated) Secondary Antibody - Background

Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency.