

Anti-MOUSE IgG (H&L) (Peroxidase Conjugated) Secondary Antibody
Rabbit Polyclonal, Peroxidase (Horseradish)
Catalog # ASR1935**Specification****Anti-MOUSE IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Product Information**

Description	Anti-MOUSE IgG (H&L) (RABBIT) Antibody Peroxidase Conjugated
Host	Rabbit
Conjugate	Peroxidase (Horseradish)
Target Species	Mouse
Clonality	Polyclonal
Application	WB, E, IC
Application Note	ELISA 1:50,000-1:300,000;Western Blot 1:1,000-1:5,000;Immunochemistry 1:500-1:2,500
Physical State	Lyophilized
Host Isotype	IgG
Target Isotype	IgG (H&L)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Mouse IgG whole molecule
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Anti-MOUSE IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Additional Information**Shipping Condition**

Ambient

Purity

HRP Secondary Antibody Conjugate was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Mouse IgG and Mouse Serum.

Storage Condition

Store mouse second antibody conjugate at 4° C prior to restoration. For extended storage aliquot antibody and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.


Anti-MOUSE IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Protein Information

Anti-MOUSE IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-MOUSE IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Images



Western Blot of Abcepta's Anti-Mouse IgG Antibody Peroxidase Conjugated detecting mouse IgG. Mouse IgG (1.0 mg) was separated on a 4-20% gradient gel and then transferred to Whatman Protran BA 85 nitrocellulose (0.45 μ m pore size), and blocked with 5% BLOTTO o/n at 4° C. The membrane was then probed with a 1:2,500 dilution of HRP Rb-a-Mouse IgG [H&L] for 45 min at RT. FemtoMax Super Sensitive Chemiluminescent HRP Substrate . Was added and the image was captured using a BioSpectrum Imaging System with a BioChem 500 -28C cooled 4.0 Megapixel CCD Camera from UVP (www.uvp.com, Upland, CA). Binning: 5X5. Exposure time: 2 min. Aperture open. No emission or excitation filters used.

Anti-MOUSE IgG (H&L) (Peroxidase Conjugated) Secondary Antibody - Background

Mouse secondary antibody conjugated to horseradish peroxidase is available in a variety of formats. Anti-IgG Secondary Antibody conjugate is Suitable for western blot, ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring lot-to-lot consistency.