

Anti-Human IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody
Goat Polyclonal, Peroxidase (Horseradish)
Catalog # ASR2047

Specification

Anti-Human IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Product Information

Description	Anti-HUMAN IgG (H&L) (GOAT) Antibody Peroxidase Conjugated (Min X MOUSE Serum Proteins)
Host Conjugate	Goat Peroxidase (Horseradish)
Target Species	Human
Reactivity	Human
Clonality	Polyclonal
Application	,1,10,15, ELISA
Application Note	1:250,000;Western Blot 1:2,000-1:20,000;Immunochemistry 1:500-1:2,500
Physical State	Lyophilized
Host Isotype	IgG
Target Isotype	IgG (H&L)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Anti-Human IgG whole molecule was produced by repeated immunization with Human IgG whole molecule in goat.
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin

Anti-Human IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Background

Anti-Human IgG whole molecule antibody generated in goat detects specifically human IgG whole molecule. This secondary antibody anti-Human is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot, immunoprecipitation and more generally immunoassays.

Preservative

**and Protease free
0.01% (w/v)
Gentamicin
Sulfate. Do NOT
add Sodium
Azide!**

**Anti-Human IgG (H&L) (Peroxidase Conjugated)
Pre-Adsorbed Secondary Antibody - Additional
Information**

Shipping Condition

Ambient

Purity

Human IgG (H&L) Antibody Peroxidase Conjugated was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Human IgG and Human Serum. No reaction was observed against Mouse Serum Proteins.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

**Anti-Human IgG (H&L) (Peroxidase Conjugated)
Pre-Adsorbed Secondary Antibody - Protein
Information**

Anti-Human IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Human IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Citations

- [Anti-CII antibody as a novel indicator to assess disease activity in systemic lupus erythematosus.](#)