

**F(ab')<sub>2</sub> Anti-Chicken IgG (H&L) (Rhodamine Conjugated) Secondary Antibody**  
**Rabbit Polyclonal, Rhodamine (TRITC)**  
**Catalog # ASR2070****Specification**

---

**F(ab')<sub>2</sub> Anti-Chicken IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Product Information**

Description	<b>F(ab')<sub>2</sub> Anti-CHICKEN IgG [H&amp;L] (RABBIT) Antibody Rhodamine Conjugated</b>
Host	<b>Rabbit</b>
Conjugate	<b>Rhodamine (TRITC)</b>
Target Species	<b>Chicken</b>
Clonality	<b>Polyclonal</b>
Physical State	<b>Lyophilized</b>
Host Isotype	<b>IgG F(ab')<sub>2</sub></b>
Target Isotype	<b>IgG (H&amp;L)</b>
Buffer	<b>0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2</b>
Immunogen	<b>Chicken IgG whole molecule</b>
Reconstitution Volume	<b>500 µL</b>
Reconstitution Buffer	<b>Restore with deionized water (or equivalent)</b>

**F(ab')<sub>2</sub> Anti-Chicken IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Additional Information****Shipping Condition**

Ambient

**Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Chicken IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Chicken IgG and Chicken Serum. No reaction was observed against anti-Pepsin or anti-Rabbit IgG F(c).

**Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

**F(ab')<sub>2</sub> Anti-Chicken IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Protein Information**

**F(ab')<sub>2</sub> Anti-Chicken IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**F(ab')<sub>2</sub> Anti-Chicken IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Images****F(ab')<sub>2</sub> Anti-Chicken IgG (H&L) (Rhodamine Conjugated) Secondary Antibody - Background**

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.