

**F(ab')2 Anti-Goat IgG (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody
Donkey Polyclonal, Rhodamine (TRITC)
Catalog # ASR2078**

Specification

**F(ab')2 Anti-Goat IgG (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody -
Product Information**

Description	F(ab')2 Anti-GOAT IgG [H&L] (DONKEY) Antibody Rhodamine Conjugated Min X Ch GP Ham Hs Hu Ms Rb & Rt Serum Proteins
Host	Donkey
Conjugate	Rhodamine (TRITC)
FP Value	2.1 moles Rhodamine (TRITC) per mole of IgG F(ab')2
Target Species	Goat
Clonality	Polyclonal
Application	IF, FC
Application Note	FLISA 1:10,000-1:50,000; IF Microscopy 1:1,000-1:5,000; Flow Cytometry 1:500-1:2,500
Physical State	Lyophilized
Host Isotype	IgG F(ab')2
Target Isotype	IgG (H&L)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Goat IgG whole molecule
Reconstitution Volume	500 µL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Sodium Azide

**F(ab')2 Anti-Goat IgG (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody -
Additional Information**

Shipping Condition

Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunolectrophoresis resulted in a single precipitin arc against anti-Donkey Serum, Goat IgG and Goat Serum. No reaction was observed against anti-Pepsin, anti-Donkey IgG F(c) or Chicken, Guinea Pig, Hamster, Horse, Human, Mouse, Rabbit and Rat Serum Proteins.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C

or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

F(ab')2 Anti-Goat IgG (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Protein Information**F(ab')2 Anti-Goat IgG (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

F(ab')2 Anti-Goat IgG (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Images**F(ab')2 Anti-Goat IgG (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Background**

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.