

## F(ab')2 Anti-Mouse IgG (H&L) (Biotin Conjugated) Secondary Antibody

Goat Polyclonal, Biotin Catalog # ASR2090

### **Specification**

# F(ab')2 Anti-Mouse IgG (H&L) (Biotin Conjugated) Secondary Antibody - Product Information

Description F(ab')2 Anti-MOUSE IgG (H&L)
(GOAT) Antibody Biotin Conjugated

Host Goat
Conjugate Biotin
Target Species Mouse
Clonality Polyclonal
Application WB, E, IC

Application Note ELISA 1:20,000-1:100,000; Western Blot

1:2,000-1:10,000;Immunochemistry

Physical State
Host Isotype
Target Isotype

1:1,000-1:5,000
Lyophilized
IgG F(ab')2
IgG (H&L)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Mouse IgG whole molecule

Reconstitution Volume 500 µL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

## F(ab')2 Anti-Mouse IgG (H&L) (Biotin Conjugated) Secondary Antibody - Additional Information

### **Shipping Condition**

**Ambient** 

**Immunogen** 

#### **Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Goat Serum, Mouse IgG and Mouse Serum. No reaction was observed against anti-Pepsin or anti-Goat IgG F(c).

#### **Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

#### **Precautions Note**



This product is for research use only and is not intended for therapeutic or diagnostic applications.

# F(ab')2 Anti-Mouse IgG (H&L) (Biotin Conjugated) Secondary Antibody - Protein Information

### F(ab')2 Anti-Mouse IgG (H&L) (Biotin Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

F(ab')2 Anti-Mouse IgG (H&L) (Biotin Conjugated) Secondary Antibody - Images

F(ab')2 Anti-Mouse IgG (H&L) (Biotin Conjugated) Secondary Antibody - Background

Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency.