

# F(ab')2 Anti-Sheep IgG (H&L) Pre-Adsorbed Secondary Antibody

Donkey Polyclonal, Unconjugated Catalog # ASR2104

#### **Specification**

#### F(ab')2 Anti-Sheep IgG (H&L) Pre-Adsorbed Secondary Antibody - Product Information

Description F(ab')2 Anti-SHEEP IgG [H&L]

(DONKEY) Antibody Min X Ch GP Ham Hs

**Hu Ms Rb & Rt Serum Proteins** 

Host **Donkey** 

Conjugate Unconjugated

Target Species Sheep Clonality Polyclonal Application WB, IHC, E

Application Note ELISA 1:20,000-1:100,000; Western Blot

1:2,000-1:10,000;Immunohistochemistry

1:1,000-1:5,000

Physical State Liquid (sterile filtered)

Host Isotype IgG F(ab')2
Target Isotype IgG (H&L)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Immunogen Sheep IgG whole molecule

Stabilizer None

Preservative 0.01% (w/v) Sodium Azide

## F(ab')2 Anti-Sheep IgG (H&L) Pre-Adsorbed Secondary Antibody - Additional Information

### **Shipping Condition**

Wet Ice

#### **Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Sheep IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Donkey Serum, Sheep IgG and Sheep Serum. No reaction was observed against anti-Pepsin, anti-Donkey IgG F(c) or Chicken, Guinea Pig, Hamster, Horse, Human, Mouse, Rabbit and Rat Serum Proteins.

#### **Storage Condition**

Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

#### **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

### F(ab')2 Anti-Sheep IgG (H&L) Pre-Adsorbed Secondary Antibody - Protein Information



#### F(ab')2 Anti-Sheep IgG (H&L) Pre-Adsorbed Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# F(ab')2 Anti-Sheep IgG (H&L) Pre-Adsorbed Secondary Antibody - Images

#### F(ab')2 Anti-Sheep IgG (H&L) Pre-Adsorbed Secondary Antibody - Background

F(ab')2 Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab)2 fragments penetrate into tissue samples and show better antigen recognition and signal generation in IHC. F(ab)2 fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab')2 Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.