

Anti-Ferret IgA (Biotin Conjugated) Secondary Antibody
Goat Polyclonal, Biotin
Catalog # ASR2121**Specification****Anti-Ferret IgA (Biotin Conjugated) Secondary Antibody - Product Information**

Description	Anti-FERRET IgA(GOAT) Antibody Biotin Conjugated
Host	Goat
Conjugate	Biotin
Target Species	Ferret
Clonality	Polyclonal
Application	,1,10,15,
Application Note	ELISA 1:20,000-1:100,000;Western Blot 1:2,000-1:10,000;Immunochemistry 1:1,000-1:5,000
Physical State	Lyophilized
Host Isotype	IgG
Target Isotype	IgA
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Ferret IgA alpha heavy chain
Species of Origin	Ferret
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Sodium Azide

Anti-Ferret IgA (Biotin Conjugated) Secondary Antibody - Additional Information**Shipping Condition**

Ambient

Purity

Anti-FERRET IgA (alpha chain) (GOAT) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Ferret IgA coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Ferret IgA and Ferret Serum. No reaction was observed against Ferret IgM or Ferret IgG.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Ferret IgA (Biotin Conjugated) Secondary Antibody - Protein Information

Anti-Ferret IgA (Biotin Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Ferret IgA (Biotin Conjugated) Secondary Antibody - Images

Anti-Ferret IgA (Biotin Conjugated) Secondary Antibody - Background

Ferret IgA whole molecule antibody generated in goat detects specifically ferret IgA whole molecule. Anti-Human IgA is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot, immunoprecipitation and more generally immunoassays