

Anti-Ferret IgG (Biotin Conjugated) Secondary Antibody
Goat Polyclonal, Biotin
Catalog # ASR2123**Specification****Anti-Ferret IgG (Biotin Conjugated) Secondary Antibody - Product Information**

Description	Anti-FERRET IgG (GOAT) Antibody Biotin Conjugated
Host	Goat
Conjugate	Biotin
Target Species	Ferret
Clonality	Polyclonal
Application	,1,10,15,
Application Note	ELISA 1:20,000-1:100,000;Western Blot 1:2,000-1:10,000;Immunochemistry 1:1,000-1:5,000
Physical State	Lyophilized
Host Isotype	IgG
Target Isotype	IgG (gamma chain)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Ferret IgG gamma heavy chain
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Sodium Azide

Anti-Ferret IgG (Biotin Conjugated) Secondary Antibody - Additional Information**Shipping Condition**

Ambient

Purity

Anti-FERRET IgG (gamma chain) (GOAT) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Ferret IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Anti-FERRET IgG (gamma chain) (GOAT) Antibody assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Ferret IgG and Ferret Serum. No reaction was observed against Ferret IgA or Ferret IgM.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Ferret IgG (Biotin Conjugated) Secondary Antibody - Protein Information

Anti-Ferret IgG (Biotin Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Ferret IgG (Biotin Conjugated) Secondary Antibody - Images

Anti-Ferret IgG (Biotin Conjugated) Secondary Antibody - Background

Anti-FERRET IgG (gamma chain) (GOAT) Antibody generated in goat detects specifically Ferret IgG gamma chain. Ferret antibody is ideal for investigators involved in Serum Protein Component research.