

Anti-Ferret IgM (mu chain) (Texas Red[™] Conjugated) Secondary Antibody Goat Polyclonal, Texas Red[®]

Catalog # ASR2126

Specification

Anti-Ferret IgM (mu chain) (Texas Red[™] Conjugated) Secondary Antibody - Product Information

Description Host Conjugate FP Value Target Species Clonality Application Application Note	Anti-FERRET IgM (mu chain) (GOAT) Antibody Texas Red™ Conjugated Goat Texas Red® 3.4 moles Texas Red® per mole of IgG Ferret Polyclonal IF, FC FLISA 1:10,000-1:50,000;IF Microscopy 1:1,000-1:5,000;FlowCytometry 1:500-1:2,500
Physical State Host Isotype Target Isotype Buffer	Lyophilized IgG IgM μ chain 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Ferret IgM mu heavy chain
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Sodium Azide

Anti-Ferret IgM (mu chain) (Texas Red[™] Conjugated) Secondary Antibody - Additional Information

Shipping Condition Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Ferret IgM and Ferret Serum. No reaction was observed against Ferret IgG.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note



This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Ferret IgM (mu chain) (Texas Red[™] Conjugated) Secondary Antibody - Protein Information

Anti-Ferret IgM (mu chain) (Texas Red[™] Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Ferret IgM (mu chain) (Texas Red[™] Conjugated) Secondary Antibody - Images

Anti-Ferret IgM (mu chain) (Texas Red[™] Conjugated) Secondary Antibody - Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.