

Anti-Human IgG F(c) (Biotin Conjugated) Secondary Antibody
Rabbit Polyclonal, Biotin
Catalog # ASR2185**Specification****Anti-Human IgG F(c) (Biotin Conjugated) Secondary Antibody - Product Information**

Description	Anti-HUMAN IgG F(c) (RABBIT) Antibody
Host	Biotin Conjugated
Conjugate	Rabbit
Target Species	Biotin
Reactivity	Human
Clonality	Human
Application	Polyclonal
Application Note	WB, E, IC
Physical State	ELISA 1:20,000-1:100,000;Western Blot
Host Isotype	1:2,000-1:10,000;Immunochemistry
Target Isotype	1:1,000-1:5,000
Buffer	Lyophilized
Immunogen	IgG
Reconstitution Volume	IgG F(c)
Reconstitution Buffer	0.02 M Potassium Phosphate, 0.15 M
Stabilizer	Sodium Chloride, pH 7.2
Preservative	Anti-Human IgG F(c) was produced by
	repeated immunization with human IgG
	F(c) fragment in rabbit.
	1.0 mL
	Restore with deionized water (or
	equivalent)
	10 mg/mL Bovine Serum Albumin (BSA) -
	Immunoglobulin and Protease free
	0.01% (w/v) Sodium Azide

Anti-Human IgG F(c) (Biotin Conjugated) Secondary Antibody - Additional Information**Shipping Condition**

Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Rabbit Serum, Human IgG, Human IgG F(c) and Human Serum. No reaction was observed against Human IgG F(ab')₂.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Human IgG F(c) (Biotin Conjugated) Secondary Antibody - Protein Information**Anti-Human IgG F(c) (Biotin Conjugated) Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Human IgG F(c) (Biotin Conjugated) Secondary Antibody - Images**Anti-Human IgG F(c) (Biotin Conjugated) Secondary Antibody - Background**

Anti-Human F(c) biotin conjugated antibody generated in rabbit detects specifically human IgG F(c). This secondary biotin conjugated antibody anti-Human is ideal for investigators who routinely perform titration assays, western-blot, immunoprecipitation and more generally immunoassays.