

Fab Anti-Goat IgG (Biotin Conjugated) Secondary Antibody
Donkey Polyclonal, Biotin
Catalog # ASR2286

Specification

Fab Anti-Goat IgG (Biotin Conjugated) Secondary Antibody - Product Information

Description	F(ab) Anti-GOAT IgG (H&L) (DONKEY) Antibody Biotin Conjugated
Host	Donkey
Conjugate	Biotin
FP Value	10-20 moles Biotin per mole of donkey IgG
Target Species	F(ab)
Clonality	Goat
Application	Polyclonal
Application Note	WB, E, IC ELISA 1:600,000;Western Blot 1:5,000-1:20,000;Immunochemistry 1:1,000-1:5,000
Physical State	Lyophilized
Host Isotype	IgG F(ab)
Target Isotype	IgG (H&L)
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Goat IgG whole molecule
Species of Origin	donkey
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Sodium Azide

Fab Anti-Goat IgG (Biotin Conjugated) Secondary Antibody - Additional Information

Shipping Condition

Ambient

Purity

F(ab) Anti-Goat IgG (H&L) was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin and anti-Donkey Serum. No reaction was observed against anti-Papain or anti-Donkey IgG F(c).

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Fab Anti-Goat IgG (Biotin Conjugated) Secondary Antibody - Protein Information**Fab Anti-Goat IgG (Biotin Conjugated) Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Fab Anti-Goat IgG (Biotin Conjugated) Secondary Antibody - Images**Fab Anti-Goat IgG (Biotin Conjugated) Secondary Antibody - Background**

F(ab) Goat IgG antibody is polyclonal that reacts with the heavy chains of IgG and the light chains common to most immunoglobulins from goat. No cross-reactivity has been detected against non-immunoglobulin serum proteins. F(ab) Anti-Goat IgG antibody is ideal for investigators involved in serum component research.