

Anti-Mouse IgG IgA IgM (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody
Goat Polyclonal, Rhodamine (TRITC)
Catalog # ASR2344

Specification

Anti-Mouse IgG IgA IgM (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Product Information

| | |
|-----------------------|--|
| Description | Anti-MOUSE IgG IgA IgM (H&L) (GOAT) Antibody Rhodamine Conjugated (Min X Bv, Hs, Hu Serum Proteins) |
| Host | Goat |
| Conjugate | Rhodamine (TRITC) |
| FP Value | 2.9 moles Rhodamine (TRITC) per mole of IgG |
| Target Species | Mouse |
| Reactivity | Mouse |
| Clonality | Polyclonal |
| Application | IF, FC |
| Application Note | FLISA 1:10,000-1:50,000;IF Microscopy 1:1,000-1:5,000;FlowCytometry 1:500-1:2,500 |
| Physical State | Lyophilized |
| Host Isotype | IgG |
| Target Isotype | IgG IgA IgM |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Immunogen | Mouse IgG IgA and IgM whole molecule |
| Reconstitution Volume | 1.0 mL |
| Reconstitution Buffer | Restore with deionized water (or equivalent) |
| Stabilizer | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |
| Preservative | 0.01% (w/v) Sodium Azide |

Anti-Mouse IgG IgA IgM (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Additional Information

Shipping Condition

Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum. No reaction was observed against bovine, horse or human serum proteins. This reagent is suitable for the detection of all mouse immunoglobulin isotypes and chain combinations.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Mouse IgG IgA IgM (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Protein Information**Anti-Mouse IgG IgA IgM (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Mouse IgG IgA IgM (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Images**Anti-Mouse IgG IgA IgM (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Background**

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.