

F(ab')2 Anti-Goat IgG (H&L) (Biotin Conjugated) Pre-Adsorbed Secondary Antibody Rabbit Polyclonal, Biotin

Catalog # ASR2381

Specification

F(ab')2 Anti-Goat IgG (H&L) (Biotin Conjugated) Pre-Adsorbed Secondary Antibody - Product Information

Description F(ab')2 Anti-GOAT IgG (H&L)

(RABBIT) Antibody Biotin Conjugated Min X

Human Serum Proteins

Host Rabbit Conjugate Biotin

FP Value ~7-8 moles Biotin per mole of IgG

F(ab')2

Target Species
Clonality
Application

Goat
Polyclonal
WB, E, IC

Application Note ELISA 1:20,000-1:100,000; Western Blot

1:2,000-1:10,000;Immunochemistry

Physical State
Host Isotype
Target Isotype

1:1,000-1:5,000
Lyophilized
IgG F(ab')2
IgG (H&L)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Goat IgG whole molecule

500 μL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.05% (w/v) Sodium Azide

F(ab')2 Anti-Goat IgG (H&L) (Biotin Conjugated) Pre-Adsorbed Secondary Antibody - Additional Information

Shipping Condition

Ambient

Immunogen

Reconstitution Volume

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Rabbit Serum, Goat IgG and Goat Serum. No reaction was observed against Human Serum Proteins.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted



Tel: 858.875.1900 Fax: 858.875.1999

liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

F(ab')2 Anti-Goat IgG (H&L) (Biotin Conjugated) Pre-Adsorbed Secondary Antibody -**Protein Information**

F(ab')2 Anti-Goat IgG (H&L) (Biotin Conjugated) Pre-Adsorbed Secondary Antibody -**Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

F(ab')2 Anti-Goat IgG (H&L) (Biotin Conjugated) Pre-Adsorbed Secondary Antibody -

F(ab')2 Anti-Goat IgG (H&L) (Biotin Conjugated) Pre-Adsorbed Secondary Antibody -**Background**

Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency.