

**F(ab')₂ Anti-Goat IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody
Donkey Polyclonal, Peroxidase (Horseradish)
Catalog # ASR2382****Specification****F(ab')₂ Anti-Goat IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody -
Product Information**

Description	F(ab')₂ Anti-GOAT IgG [H&L] (DONKEY) Antibody Peroxidase Conjugated Min X Ch GP Ham Hs Hu Ms Rb & Rt Serum Proteins Donkey Peroxidase (Horseradish) Goat Polyclonal ,1,10,15, ELISA 1:10,000-1:50,000;Western Blot 1:2,000-1:10,000;Immunochemistry 1:500-1:5,000 Lyophilized IgG F(ab')₂ IgG (H&L) 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Goat IgG whole molecule 500 µL Restore with deionized water (or equivalent) 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!
Host	
Conjugate	
Target Species	
Clonality	
Application	
Application Note	
Physical State	
Host Isotype	
Target Isotype	
Buffer	
Immunogen	
Reconstitution Volume	
Reconstitution Buffer	
Stabilizer	
Preservative	

**F(ab')₂ Anti-Goat IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody -
Additional Information****Shipping Condition**

Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Donkey Serum, Goat IgG and Goat Serum. No reaction was observed against anti-Pepsin, anti-Donkey IgG F(c) or Chicken, Guinea Pig, Hamster, Horse, Human, Mouse, Rabbit and Rat Serum Proteins.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after

standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

F(ab')₂ Anti-Goat IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Protein Information**F(ab')₂ Anti-Goat IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

F(ab')₂ Anti-Goat IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Images