

F(ab')2 Anti-Human IgG F(ab')2 (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody

Goat Polyclonal, Peroxidase (Horseradish) Catalog # ASR2389

Specification

F(ab')2 Anti-Human IgG F(ab')2 (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Product Information

Description F(ab')2 Anti-HUMAN IgG

F(ab')2 (GOAT) Antibody Peroxidase Conjugated Min X Bv Hs Ms & Rt Serum

Proteins

Host Goat

Conjugate Peroxidase (Horseradish)

Target Species
Reactivity
Human
Clonality
Application
Human
Polyclonal
WB, E, IC

Application Note ELISA 1:10,000-1:40,000; Western Blot

1:1,000-1:5,000;Immunochemistry

Physical State
Host Isotype
Target Isotype

1:500-1:2,500
Lyophilized
IgG F(ab')2
IgG F(ab')2

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Immunogen Anti-Human IgG F(ab')2 fragment

antibody was produced by repeated immunization with Human IgG F(ab')2 fragment in goat.

Reconstitution Volume 500 μL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Gentamicin Sulfate. Do NOT

add Sodium Azide!

F(ab')2 Anti-Human IgG F(ab')2 (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Additional Information

Shipping Condition Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Human IgG, Human IgG F(ab')2 and Human Serum. No reaction was observed against anti-Pepsin,



anti-Goat IgG F(c), Human IgG F(c) or Bovine, Horse, Mouse and Rat Serum Proteins.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

F(ab')2 Anti-Human IgG F(ab')2 (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Protein Information

F(ab')2 Anti-Human IgG F(ab')2 (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

F(ab')2 Anti-Human IgG F(ab')2 (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Images

F(ab')2 Anti-Human IgG F(ab')2 (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Background

Anti-Human IgG F(ab')2 fragment antibody generated in goat detects specifically Human IgG F(ab')2 fragment. This secondary antibody anti-Human is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot, immunoprecipitation and more generally immunoassays.