

**Dog IgG Fab**  
**Catalog # ASR2414****Specification**

---

**Dog IgG Fab - Product Information**

Description	<b>DOG IgG F(ab) fragment</b>
Conjugate	<b>Unconjugated</b>
Physical State	<b>Lyophilized</b>
Host Isotype	<b>IgG F(ab)</b>
Buffer	<b>0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2</b>
Species of Origin	<b>Dog</b>
Reconstitution Volume	<b>1.0 mL</b>
Reconstitution Buffer	<b>Restore with deionized water (or equivalent)</b>
Stabilizer	<b>None</b>
Preservative	<b>0.01% (w/v) Sodium Azide</b>

**Dog IgG Fab - Additional Information****Shipping Condition**

Wet Ice

**Purity**

This product was prepared from normal serum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Dog IgG, anti-Dog IgG F(ab')<sub>2</sub> and anti-Dog Serum. No reaction was observed against anti-Dog IgG F(c) or anti- Papain.

**Storage Condition**

Store vial at 4° C prior to restoration. Restore with 1.0 mL of deionized water (or equivalent). For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Precautions Note**


This product is for research use only and is not intended for therapeutic or diagnostic applications.

**Dog IgG Fab - Protein Information****Dog IgG Fab - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Dog IgG Fab - Images



SDS-Page of Dog IgG F(ab) Fragment. Lane 1: Dog F(ab) - Reduced. Lane 2: Dog F(ab) - Non-reduced. Load: 1.0  $\mu$ g per lane. Predicted/Observed size: 25 kDa - Reduced, 50 kDa - Non-reduced for F(ab) fragment. Other band(s): None.