

Goat IgG F(ab')2 (Agarose Conjugated)

Catalog # ASR2415

Specification

Goat IgG F(ab')2 (Agarose Conjugated) - Product Information

Description GOAT IgG F(ab')2 fragment Agarose

> Conjugated Unconjugated

Conjugate

Physical State Suspension of agarose beads Buffer

0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2

Species of Origin Goat Stabilizer None

Preservative 0.01% (w/v) Sodium Azide

Goat IgG F(ab')2 (Agarose Conjugated) - Additional Information

Shipping Condition

Wet Ice

Purity

This product is an purified Goat IgG isolated from normal Goat Serum digested by the enzyme pepsin and purified by chromatography coupled to activated agarose. Sufficient antibody capacity is provided to bind a minimum of 5.0 mg of anti-Goat IgG.

Storage Condition

Store vial at 4° C prior to opening. DO NOT FREEZE.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Goat IgG F(ab')2 (Agarose Conjugated) - Protein Information


Goat IgG F(ab')2 (Agarose Conjugated) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Goat IgG F(ab')2 (Agarose Conjugated) - Images





SDS-PAGE of Goat IgG F(ab')2 Fragment Agarose Conjugated . Lane M: 5 μ L Opal Prestained Marker . Lane 1: Reduced Goat IgG Whole Molecule . Lane 2: Reduced Goat IgG F(c) Fragment . Lane 3: Reduced Goat IgG F(ab')2 Fragment Agarose Conjugated . Lane 4: Reduced Goat IgM Whole Molecule . Load: 1 μ g for IgG, F(c) and F(ab')2; 3 μ g for IgM. Predicted/Observed size: IgG at 50 and 25 kDa; F(c) at 25 kDa; F(ab) at 25 kDa; IgM at 70 and 23 kDa. Observed F(c) Fragment migrates slightly higher.