

**Rabbit IgG F(ab')<sub>2</sub> (Agarose Conjugated)**  
**Catalog # ASR2425****Specification**

---

**Rabbit IgG F(ab')<sub>2</sub> (Agarose Conjugated) - Product Information**

Description	<b>RABBIT IgG F(ab')<sub>2</sub> fragment</b>
Conjugate	<b>Agarose Conjugated</b>
Physical State	<b>Unconjugated</b>
Buffer	<b>Suspension of agarose beads</b>
	<b>0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2</b>
Species of Origin	<b>Rabbit</b>
Stabilizer	<b>None</b>
Preservative	<b>0.5% (w/v) Sodium Azide</b>

**Rabbit IgG F(ab')<sub>2</sub> (Agarose Conjugated) - Additional Information****Shipping Condition**

Wet Ice

**Purity**

This product is an purified Rabbit IgG isolated from normal Rabbit Serum digested by the enzyme pepsin and purified by chromatography coupled to activated agarose. Sufficient antibody capacity is provided to bind a minimum of 5 mg of anti-Rabbit IgG.

**Storage Condition**

Store vial at 4° C prior to opening. DO NOT FREEZE.

**Precautions Note**


This product is for research use only and is not intended for therapeutic or diagnostic applications.

**Rabbit IgG F(ab')<sub>2</sub> (Agarose Conjugated) - Protein Information****Rabbit IgG F(ab')<sub>2</sub> (Agarose Conjugated) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Rabbit IgG F(ab')<sub>2</sub> (Agarose Conjugated) - Images**



SDS-PAGE of Rabbit IgG F(ab')<sub>2</sub> Fragment Agarose Conjugated . Lane M: 3  $\mu$ L Opal Prestained Marker . Lane 1: Reduced Rabbit IgG Whole Molecule . Lane 2: Reduced Rabbit IgG F(ab')<sub>2</sub> Fragment Agarose Conjugated . Lane 3: Reduced Rabbit IgG F(c) Fragment . Lane 4: Reduced Rabbit IgM Whole Molecule . Load: 1  $\mu$ g for F(ab')<sub>2</sub> and F(c); 1.2  $\mu$ g for IgG and IgM. Predicted/Observed size: IgG at 50 and 25 kDa; F(c) at 25 kDa; F(ab')<sub>2</sub> at 25 kDa; IgM at 70 and 23 kDa. Observed F(c) Fragment migrates slightly higher.