

Bovine IgM
Catalog # ASR2551**Specification****Bovine IgM - Product Information**

Description	BOVINE IgM whole molecule
Conjugate	Unconjugated
Physical State	Liquid (sterile filtered)
Host Isotype	IgM
Buffer	0.1 M Tris Chloride, 0.5 M Sodium Chloride, pH 8.0
Species of Origin	Bovine
Preservative	0.1% (w/v) Sodium Azide

Bovine IgM - Additional Information**Shipping Condition**

Wet Ice

Purity

Bovine IgM whole molecule was prepared from normal serum by a multi-step process which includes delipidation, selective precipitation and tandem molecular sieve chromatography followed by extensive dialysis against the buffer stated above. Bovine IgM whole molecule assayed by immunoelectrophoresis resulted in a single precipitin arc against anti-Bovine Serum and anti-Bovine IgM (μ chain specific). No reaction was observed against anti-Bovine IgG F(c). Some light chain cross reactivity will occur with anti-Bovine IgG.

Storage Condition

Store vial at 4° C prior to opening. Bovine IgM whole molecule is stable 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage mix with an equal volume of glycerol, aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Bovine IgM - Protein Information**Bovine IgM - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Bovine IgM - Images**Bovine IgM - Background**

Immunoglobulin M is the largest antibody isotype and the first to be secreted against an initial exposure to antigen. IgM is predominantly produced in the spleen. Formed from covalently linking 5 immunoglobulins together, the approximate molecular weight of IgM is 900kDa and possesses 10 binding sites (though due to the size of most antigens, not all sites are capable of binding at once). Due to this large size, IgM is typically isolated to the serum. Bovine IgM whole molecule is ideal for investigators in Immunology, Microbiology, and Cell Biology.