

# Anti-Mouse IgG F(c) (Peroxidase Conjugated) Secondary Antibody

Rabbit Polyclonal, Peroxidase (Horseradish) Catalog # ASR2753

#### Specification

# Anti-Mouse IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Product Information

Description

- Host Conjugate Target Species Clonality Application Application Note
- Physical State Host Isotype Target Isotype Buffer

Immunogen Reconstitution Volume Reconstitution Buffer

Stabilizer

Preservative

Anti-MOUSE IgG F(c) (RABBIT) Antibody **Peroxidase Conjugated** Rabbit Peroxidase (Horseradish) Mouse Polyclonal WB, E, IC ELISA 1:10,000-1:50,000;Western Blot 1:1,000-1:10,000;Immunochemistry 1:500-1:2.500 Lyophilized laG laG F(c) 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Mouse IgG F(c) fragment 1.0 mL Restore with deionized water (or equivalent) 10 mg/mL Bovine Serum Albumin (BSA) -**Immunoglobulin and Protease free** 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

# Anti-Mouse IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Additional Information

#### Shipping Condition Ambient

#### **Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Mouse IgG, Mouse IgG F(c) and Mouse Serum. No reaction was observed against Mouse IgG F(ab')2 fragment.

#### Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

#### **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.



# Anti-Mouse IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Protein Information

### Anti-Mouse IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

# Anti-Mouse IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Images