

F(ab')₂ Anti-Human IgG IgA IgM (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody**Goat Polyclonal, Rhodamine (TRITC)
Catalog # ASR2828****Specification****F(ab')₂ Anti-Human IgG IgA IgM (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Product Information**

Description	F(ab')₂ Anti-HUMAN IgG IgA IgM (H&L) (GOAT) Antibody Rhodamine Conjugated Min X MOUSE Serum Proteins Goat
Host	Rhodamine (TRITC)
Conjugate	3.9 moles Rhodamine (TRITC) per mole of Human IgG F(ab')₂
FP Value	Human
Target Species	Polyclonal
Clonality	,3,4,
Application	FLISA 1:10,000-1:50,000;IF Microscopy 1:1,000-1:5,000;FlowCytometry 1:500-1:2,500
Application Note	Lyophilized
Physical State	IgG F(ab')₂
Host Isotype	IgG IgA IgM
Target Isotype	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Buffer	Human IgG, IgA, and IgM whole molecules
Immunogen	Human
Species of Origin	1.0 mL
Reconstitution Volume	Restore with deionized water (or equivalent)
Reconstitution Buffer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Stabilizer	0.01% (w/v) Sodium Azide
Preservative	

F(ab')₂ Anti-Human IgG IgA IgM (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Additional Information**Shipping Condition**

Ambient

Purity

F(ab')₂ Anti-HUMAN IgG IgA IgM (H&L) (GOAT) Antibody was prepared from polyspecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum. No reaction was observed against anti-Goat IgG F(c), anti-Pepsin or Mouse Serum Proteins. This product is suitable for the detection of all Human immunoglobulin classes, isotypes and chain combinations.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

F(ab')₂ Anti-Human IgG IgA IgM (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Protein Information**F(ab')₂ Anti-Human IgG IgA IgM (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

F(ab')₂ Anti-Human IgG IgA IgM (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Images**F(ab')₂ Anti-Human IgG IgA IgM (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Background**

F(ab')₂ Anti-HUMAN IgG IgA IgM (H&L) (GOAT) Antibody is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.