

**Cat IgM**  
**Catalog # ASR2832****Specification****Cat IgM - Product Information**

Description	CAT IgM whole molecule
Conjugate	Unconjugated
Physical State	Liquid (sterile filtered)
Host Isotype	IgM
Buffer	0.1 M Tris Chloride, 0.5 M Sodium Chloride, pH 8.0
Species of Origin	Cat
Preservative	0.1% (w/v) Sodium Azide

**Cat IgM - Additional Information****Shipping Condition**

Wet Ice

**Purity**

Cat IgM whole molecule was prepared from normal serum by a multi-step process which includes delipidation, selective precipitation and tandem molecular sieve chromatography followed by extensive dialysis against the buffer stated above. Cat IgM whole molecule was assayed by immunoelectrophoresis resulted in a single precipitin arc against anti-Cat Serum and anti-Cat IgM ( $\mu$  chain specific). No reaction was observed against anti-Cat IgG F(c). Some light chain cross reactivity will occur with anti-Cat IgG.

**Storage Condition**

Store vial at 4° C prior to opening. Cat IgM whole molecule is stable 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage mix with an equal volume of glycerol, aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

**Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

**Cat IgM - Protein Information****Cat IgM - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **Cat IgM - Images**

### **Cat IgM - Background**

Immunoglobulin M is the largest antibody isotype and the first to be secreted against an initial exposure to antigen. IgM is predominantly produced in the spleen. Formed from covalently linking 5 immunoglobulins together, the approximate molecular weight of IgM is 900kDa and possesses 10 binding sites (though due to the size of most antigens, not all sites are capable of binding at once). Due to this large size, IgM is typically isolated to the serum.