

Dog IgM
Catalog # ASR2833**Specification****Dog IgM - Product Information**

Description	DOG IgM whole molecule
Conjugate	Unconjugated
Physical State	Liquid (sterile filtered)
Host Isotype	IgM
Buffer	0.1 M Tris Chloride, 0.5 M Sodium Chloride, pH 8.0
Species of Origin	Dog
Preservative	0.1% (w/v) Sodium Azide

Dog IgM - Additional Information**Shipping Condition**

Wet Ice

Purity

Dog IgM whole molecule was prepared from normal serum by a multi-step process which includes delipidation, selective precipitation and tandem molecular sieve chromatography followed by extensive dialysis against the buffer stated above. Dog IgM whole molecule was assayed by immunoelectrophoresis resulted in a single precipitin arc against anti-Dog Serum and anti-Dog IgM (μ chain specific). No reaction was observed against anti-Dog IgG F(c). Some light chain cross reactivity will occur with anti-Dog IgG.

Storage Condition

Store vial at 4° C prior to opening. Dog IgM whole molecule is stable 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage mix with an equal volume of glycerol, aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Dog IgM - Protein Information**Dog IgM - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Dog IgM - Images**Dog IgM - Background**

Immunoglobulin M is the largest antibody isotype and the first to be secreted against an initial exposure to antigen. IgM is predominantly produced in the spleen. Formed from covalently linking 5 immunoglobulins together, the approximate molecular weight of IgM is 900kDa and possesses 10 binding sites (though due to the size of most antigens, not all sites are capable of binding at once). Due to this large size, IgM is typically isolated to the serum.