

**Anti-Dog IgG F(c) (Peroxidase Conjugated) Secondary Antibody**  
**Rabbit Polyclonal, Peroxidase (Horseradish)**  
**Catalog # ASR2842****Specification****Anti-Dog IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Product Information**

|                       |  |
|-----------------------|--|
| Description           | <b>Anti-DOG IgG F(c) (RABBIT) Antibody</b><br><b>Peroxidase Conjugated</b>                                     |
| Host                  | <b>Rabbit</b>  |
| Conjugate             | <b>Peroxidase (Horseradish)</b>  |
| Target Species        | <b>Dog</b>   |
| Clonality             | <b>Polyclonal</b>  |
| Application           | <b>WB, E, IC</b>   |
| Application Note      | <b>ELISA 1:10,000-1:50,000;Western Blot</b><br><b>1:1,000-1:10,000;Immunochemistry</b><br><b>1:500-1:2,500</b> |
| Physical State        | <b>Lyophilized</b>   |
| Host Isotype          | <b>IgG</b>   |
| Target Isotype        | <b>IgG F(c)</b>  |
| Buffer                | <b>0.02 M Potassium Phosphate, 0.15 M</b><br><b>Sodium Chloride, pH 7.2</b>                                    |
| Immunogen             | <b>Dog IgG F(c) fragment</b>   |
| Reconstitution Volume | <b>1.0 mL</b>  |
| Reconstitution Buffer | <b>Restore with deionized water (or</b><br><b>equivalent)</b>  |
| Stabilizer            | <b>10 mg/mL Bovine Serum Albumin (BSA) -</b><br><b>Immunoglobulin and Protease free</b>                        |
| Preservative          | <b>0.01% (w/v) Gentamicin Sulfate. Do NOT</b><br><b>add Sodium Azide!</b>                                      |

**Anti-Dog IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Additional Information****Shipping Condition**

Ambient

**Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Dog IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Dog IgG, Dog IgG F(c) and Dog Serum. No reaction was observed against Dog IgG F(ab')<sub>2</sub>.

**Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

## **Anti-Dog IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Protein Information**

## **Anti-Dog IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **Anti-Dog IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Images**