

F(ab')2 Anti-Rat IgG (H&L) (Phycoerythrin Conjugated) Secondary Antibody

Rabbit Polyclonal, R-Phycoerythrin (RPE)
Catalog # ASR2877

Specification

Host

Conjugate

Clonality

Buffer

Target Species

Physical State

Target Isotype

Host Isotype

Immunogen

Stabilizer

Preservative

Information

Reconstitution Volume

Reconstitution Buffer

F(ab')2 Anti-Rat IgG (H&L) (Phycoerythrin Conjugated) Secondary Antibody - Product Information

Description F(ab')2 Anti-RAT IgG [H&L] (RABBIT)

Antibody Phycoerythrin conjugated

Rabbit

R-Phycoerythrin (RPE)

Rat

Polyclonal Lyophilized IgG F(ab')2

IgG (H&L)

0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2 Rat IgG whole molecule

1.0 mL

Restore with deionized water (or

equivalent)

10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

0.01% (w/v) Sodium Azide

F(ab')2 Anti-Rat IgG (H&L) (Phycoerythrin Conjugated) Secondary Antibody - Additional

Shipping Condition

Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by pepsin digestion and chromatographic separation. Coupling to R-PE was followed by size exclusion chromatography to purify conjugate from unreacted R-PE and antibody. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Phycoerythrin, anti-Rabbit Serum, Rat IgG and Rat Serum. No reaction was observed against anti-Pepsin or anti-Rabbit IgG F(c).

Storage Condition

Store vial at 4° C prior to opening. Dilute only prior to immediate use. Do not freeze after reconstitution. Store reagent in the dark. This product is stable at 4° C as an undiluted liquid. Use subdued lighting during handling and incubation of cells prior to analysis.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

F(ab')2 Anti-Rat IgG (H&L) (Phycoerythrin Conjugated) Secondary Antibody - Protein



Information

F(ab')2 Anti-Rat IgG (H&L) (Phycoerythrin Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

F(ab')2 Anti-Rat IgG (H&L) (Phycoerythrin Conjugated) Secondary Antibody - Images