

**F(ab')₂ Anti-Fragment Of Hamster IgG (H&L) (Biotin Conjugated) Pre-Adsorbed
Secondary Antibody
Goat Polyclonal, Biotin
Catalog # ASR2878**

Specification

**F(ab')₂ Anti-Fragment Of Hamster IgG (H&L) (Biotin Conjugated) Pre-Adsorbed
Secondary Antibody - Product Information**

| | |
|-----------------------|--|
| Description | F(ab')₂ Fragment of Affinity Purified Anti-GOLDEN SYRIAN & ARMENIAN HAMSTER IgG (H&L) (GOAT) Antibody Biotin Conjugated (Min X MOUSE and RAT Serum Proteins) |
| Host | Goat |
| Conjugate | Biotin |
| Target Species | Armenian and Golden Syrian Hamster |
| Clonality | Polyclonal |
| Application | WB, IHC, E |
| Application Note | ELISA 1:5000-1:25,000;Western Blot 1:2,000-1:10,000;Immunohistochemistry 1:300-1:2000 |
| Physical State | Lyophilized |
| Host Isotype | IgG F(ab')₂ |
| Target Isotype | IgG (H&L) |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Immunogen | Golden Syrian and Armenian Hamster IgG whole molecules |
| Reconstitution Volume | 500 µL |
| Reconstitution Buffer | Restore with deionized water (or equivalent) |
| Stabilizer | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |
| Preservative | 0.01% (w/v) Sodium Azide |

**F(ab')₂ Anti-Fragment Of Hamster IgG (H&L) (Biotin Conjugated) Pre-Adsorbed
Secondary Antibody - Additional Information**

Shipping Condition

Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Golden Syrian and Armenian Hamster IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Golden Syrian and Armenian Hamster IgG and Golden Syrian and Armenian Hamster Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c) and Mouse or Rat Serum Proteins.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

F(ab')₂ Anti-Fragment Of Hamster IgG (H&L) (Biotin Conjugated) Pre-Adsorbed Secondary Antibody - Protein Information**F(ab')₂ Anti-Fragment Of Hamster IgG (H&L) (Biotin Conjugated) Pre-Adsorbed Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

F(ab')₂ Anti-Fragment Of Hamster IgG (H&L) (Biotin Conjugated) Pre-Adsorbed Secondary Antibody - Images**F(ab')₂ Anti-Fragment Of Hamster IgG (H&L) (Biotin Conjugated) Pre-Adsorbed Secondary Antibody - Background**

Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency.