

Anti-Dog IgM mu chain Secondary Antibody
Rabbit Polyclonal, Unconjugated
Catalog # ASR2909**Specification**

Anti-Dog IgM mu chain Secondary Antibody - Product Information

| | |
|-----------------------|--|
| Description | Anti-DOG IgM mu chain (RABBIT) Antibody |
| Host | Rabbit |
| Conjugate | Unconjugated |
| Target Species | Dog |
| Clonality | Polyclonal |
| Application | WB, IHC, E |
| Application Note | ELISA 1:20,000-1:100,000;Western Blot 1:2,000-1:10,000;Immunohistochemistry 1:1,000-1:5,000 |
| Physical State | Lyophilized |
| Host Isotype | IgG |
| Target Isotype | IgM μ chain |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Immunogen | Dog IgM mu heavy chain |
| Reconstitution Volume | 5.0 mL |
| Reconstitution Buffer | Restore with deionized water (or equivalent) |
| Stabilizer | None |
| Preservative | 0.01% (w/v) Sodium Azide |

Anti-Dog IgM mu chain Secondary Antibody - Additional Information**Shipping Condition**

Ambient

Purity

This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Dog IgM and Dog Serum. No reaction was observed against Dog IgG.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Dog IgM mu chain Secondary Antibody - Protein Information

Anti-Dog IgM mu chain Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Dog IgM mu chain Secondary Antibody - Images