

**Anti-Guinea Pig IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody  
Donkey Polyclonal, Peroxidase (Horseradish)  
Catalog # ASR2948****Specification****Anti-Guinea Pig IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody -  
Product Information**

Description	<b>Anti-GUINEA PIG IgG (H&amp;L) (DONKEY) Antibody Peroxidase Conjugated (Min X Bv Ch Gt Ham Hs Hu Ms Rb Rt &amp; Sh Serum Proteins)</b>
Host	<b>Donkey</b>
Conjugate	<b>Peroxidase (Horseradish)</b>
Target Species	<b>Guinea Pig</b>
Reactivity	<b>Guinea Pig</b>
Clonality	<b>Polyclonal</b>
Application	<b>WB, E</b>
Application Note	<b>ELISA 1:10,000-1:50,000;Western Blot 1:1,000-1:5,000</b>
Physical State	<b>Lyophilized</b>
Host Isotype	<b>IgG</b>
Target Isotype	<b>IgG (H&amp;L)</b>
Buffer	<b>0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2</b>
Immunogen	<b>Guinea Pig IgG whole molecule</b>
Reconstitution Volume	<b>1.0 mL</b>
Reconstitution Buffer	<b>Restore with deionized water (or equivalent)</b>
Stabilizer	<b>10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free</b>
Preservative	<b>0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!</b>

**Anti-Guinea Pig IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody -  
Additional Information****Shipping Condition**

Ambient

**Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Guinea Pig IgG coupled to agarose beads followed by conjugation to fluorochrome and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Donkey Serum, Guinea Pig IgG and Guinea Pig Serum. No reaction was observed against Bovine, Chicken, Goat, Hamster, Horse, Human, Mouse, Rabbit, Rat and Sheep Serum Proteins. This antibody will react with heavy chains of Guinea Pig IgG and with light chains of most Guinea Pig immunoglobulins.

**Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C

or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

**Anti-Guinea Pig IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Protein Information****Anti-Guinea Pig IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-Guinea Pig IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody - Images**