

# Anti-Mouse IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody

Donkey Polyclonal, Peroxidase (Horseradish) Catalog # ASR2960

## **Specification**

## Anti-Mouse IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody -Product Information

Description	Anti-MOUSE IgG (H&L) (DONKEY) Antibody Peroxidase Conjugated (Min X Bv Ch Gt GP Ham Hs Hu Rb Rt & Sh Serum Proteins)
Host	Donkey
Conjugate	Peroxidase (Horseradish)
Target Species	Mouse
Reactivity	Mouse
Clonality	Polyclonal
Application	WB, E, IC
Application Note	ELISA 1:180,000;Western Blot
	1:2,000-1:20,000;Immunochemistry
	1:1,000-1:5,000
Physical State	Lyophilized
Host Isotype	lgG
Target Isotype	IgG (H&L)
Buffer	0.02 M Potassium Phosphate, 0.15 M
	Sodium Chloride, pH 7.2
Immunogen	Anti-Mouse IgG whole molecule was
	produced by repeated immunization with
	Mouse IgG whole molecule in donkey.
Reconstitution Volume	1.0 mL
Reconstitution Buffer	Restore with deionized water (or
	equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) -
	Immunoglobulin and Protease free
Preservative	0.01% (w/v) Gentamicin Sulfate. Do NOT
	add Sodium Azide!

## Anti-Mouse IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody -Additional Information

Shipping Condition Ambient

**Purity** 

Mouse IgG (H&L) Antibody Peroxidase Conjugated was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Donkey Serum, Mouse IgG and Mouse Serum. No reaction was observed against Bovine, Chicken, Goat, Guinea Pig, Hamster, Horse, Human, Rabbit, Rat and Sheep Serum Proteins

**Storage Condition** 



Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

#### **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

## Anti-Mouse IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody -Protein Information

## Anti-Mouse IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody -Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Mouse IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody -Images

kDa 150	
100 - 🕳	
80 - 🕳	
60 - 🖝	
50 -	
40 -	
30 - 🖛 🖛 ┥	
20 -	

Western Blot of Peroxidase Donkey Anti-Mouse IgG (Pre-Adsorbed) secondary antibody. Lane 1: Mouse IgG. Lane 2: None. Load: 50 ng per lane. Primary antibody: None. Secondary antibody: Peroxidase donkey secondary antibody at 1:1,000 for 60 min at RT. Blocking: MB-070 for 30 min at RT. Predicted/Observed size: 25 & 55 kDa, 25 & 55 kDa for Mouse IgG. Other band(s): None.

## Anti-Mouse IgG (H&L) (Peroxidase Conjugated) Pre-Adsorbed Secondary Antibody -Background

Anti-Mouse IgG whole molecule antibody generated in donkey detects specifically Mouse IgG whole molecule . This secondary antibody anti-Mouse is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot, immunoprecipitation and more



generally immunoassays.