

F(ab')₂ Anti-Cat IgG (H&L) (Phycoerythrin Conjugated) Secondary Antibody
Rabbit Polyclonal, R-Phycoerythrin (RPE)
Catalog # ASR2980**Specification****F(ab')₂ Anti-Cat IgG (H&L) (Phycoerythrin Conjugated) Secondary Antibody - Product Information**

| | |
|-----------------------|---|
| Description | F(ab')₂ Anti-CAT IgG [H&L] (RABBIT) Antibody Phycoerythrin conjugated |
| Host | Rabbit |
| Conjugate | R-Phycoerythrin (RPE) |
| Target Species | Cat |
| Clonality | Polyclonal |
| Physical State | Lyophilized |
| Host Isotype | IgG F(ab')₂ |
| Target Isotype | IgG (H&L) |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Immunogen | Cat IgG whole molecule |
| Reconstitution Volume | 1.0 mL |
| Reconstitution Buffer | Restore with deionized water (or equivalent) |

F(ab')₂ Anti-Cat IgG (H&L) (Phycoerythrin Conjugated) Secondary Antibody - Additional Information**Shipping Condition**

Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Cat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Cat IgG and Cat Serum. No reaction was observed against anti-Pepsin or anti-Rabbit IgG F(c).

Storage Condition

Store vial at 4° C prior to opening. Dilute only prior to immediate use. Do not freeze after reconstitution. Store reagent in the dark. This product is stable at 4° C as an undiluted liquid. Use subdued lighting during handling and incubation of cells prior to analysis.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

F(ab')₂ Anti-Cat IgG (H&L) (Phycoerythrin Conjugated) Secondary Antibody - Protein Information

F(ab')₂ Anti-Cat IgG (H&L) (Phycoerythrin Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

F(ab')₂ Anti-Cat IgG (H&L) (Phycoerythrin Conjugated) Secondary Antibody - Images