

F(ab')₂ Anti-MOUSE IgM (mu chain) (Peroxidase Conjugated) Secondary Antibody
Goat Polyclonal, Peroxidase (Horseradish)
Catalog # ASR2991**Specification****F(ab')₂ Anti-MOUSE IgM (mu chain) (Peroxidase Conjugated) Secondary Antibody - Product Information**

| | |
|-----------------------|--|
| Description | F(ab')₂ Anti-MOUSE IgM (mu chain) (GOAT) Antibody Peroxidase Conjugated |
| Host | Goat |
| Conjugate | Peroxidase (Horseradish) |
| Target Species | Mouse |
| Reactivity | Mouse |
| Clonality | Polyclonal |
| Application | WB, E, IC |
| Application Note | ELISA 1:100,000;Western Blot 1:2,000-1:10,000;Immunochemistry 1:500-1:2,500 |
| Physical State | Lyophilized |
| Host Isotype | IgG F(ab')₂ |
| Target Isotype | IgM μ chain |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Immunogen | F(ab')₂ Anti-Mouse IgM mu heavy chain was produced by repeated immunization with Mouse IgM mu heavy chain in goat. |
| Reconstitution Volume | 500 μL |
| Reconstitution Buffer | Restore with deionized water (or equivalent) |
| Stabilizer | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |
| Preservative | 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide! |

F(ab')₂ Anti-MOUSE IgM (mu chain) (Peroxidase Conjugated) Secondary Antibody - Additional Information**Shipping Condition**

Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgM coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Mouse IgM and Mouse Serum. No reaction was observed against anti-Pepsin or anti-Goat IgG F(c). Specificity was confirmed by ELISA at less than 1% cross reactivity against other mouse heavy or light chain isotypes.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

F(ab')₂ Anti-MOUSE IgM (mu chain) (Peroxidase Conjugated) Secondary Antibody - Protein Information**F(ab')₂ Anti-MOUSE IgM (mu chain) (Peroxidase Conjugated) Secondary Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

F(ab')₂ Anti-MOUSE IgM (mu chain) (Peroxidase Conjugated) Secondary Antibody - Images**F(ab')₂ Anti-MOUSE IgM (mu chain) (Peroxidase Conjugated) Secondary Antibody - Background**

F(ab')₂ Anti-Mouse IgM mu heavy chain generated in goat detects specifically Mouse IgM mu heavy chain. This secondary antibody anti-Mouse is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assays, western-blot, immunoprecipitation and more generally immunoassays.