

Tel: 858.875.1900 Fax: 858.875.1999

## F(ab')2 Anti-Rat IgG (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody Rabbit Polyclonal, Rhodamine (TRITC) Catalog # ASR3001

### **Specification**

Conjugate

Immunogen

### F(ab')2 Anti-Rat IgG (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody -**Product Information**

Description F(ab')2 Anti-RAT IgG (H&L) (RABBIT)

**Antibody Rhodamine Conjugated Min X By** 

Hs Hu Ms Rb & Sh Serum Proteins

Host Rabbit

FP Value 2.1 moles Rhodamine (TRITC) per mole of

IgG F(ab')2

**Rhodamine (TRITC)** 

**Target Species** Rat

Clonality **Polyclonal** Application IF, FC

**Application Note** FLISA 1:10,000-1:50,000; IF Microscopy

1:1,000-1:5,000;FlowCytometry

1:500-1:2,500 **Physical State** Lyophilized Host Isotype IgG F(ab')2 Target Isotype IgG (H&L)

Buffer 0.02 M Potassium Phosphate, 0.15 M

> Sodium Chloride, pH 7.2 Rat IgG whole molecule

Reconstitution Volume 500 uL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

0.01% (w/v) Sodium Azide Preservative

### F(ab')2 Anti-Rat IgG (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody -**Additional Information**

# **Shipping Condition**

**Ambient** 

### **Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Rat IgG and Rat Serum. No reaction was observed against anti-Pepsin, anti-Rabbit IgG F(c) or Bovine, Horse, Human, Mouse, Rabbit and Sheep Serum Proteins.

### **Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after



standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

#### **Precautions Note**

This product is for research use only and is not intended for therapeutic or diagnostic applications.

F(ab')2 Anti-Rat IgG (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Protein Information

F(ab')2 Anti-Rat IgG (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

F(ab')2 Anti-Rat IgG (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Images

F(ab')2 Anti-Rat IgG (H&L) (Rhodamine Conjugated) Pre-Adsorbed Secondary Antibody - Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.