

## Anti-Ferret IgM (mu chain) (Peroxidase Conjugated) Secondary Antibody

Goat Polyclonal, Peroxidase (Horseradish)
Catalog # ASR3009

### **Specification**

**Physical State** 

Host Isotype

**Immunogen** 

Reconstitution Volume

# Anti-Ferret IgM (mu chain) (Peroxidase Conjugated) Secondary Antibody - Product Information

Description Anti-FERRET IgM (mu chain) (GOAT)

Antibody Peroxidase Conjugated

Host Goat

Conjugate Peroxidase (Horseradish)

Target Species Ferret
Clonality Polyclonal
Application WB, E, IC

Application Note ELISA 1:10,000-1:50,000; Western Blot

1:1,000-1:5,000;Immunochemistry

1:500-1:2,500 Lyophilized

IgG

Target Isotype IgM μ chain

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2 Ferret IgM mu heavy chain

1.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Thimerosal

# Anti-Ferret IgM (mu chain) (Peroxidase Conjugated) Secondary Antibody - Additional Information

# **Shipping Condition**

**Ambient** 

### **Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Ferret IgM coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Ferret IgM and Ferret Serum. Specificity was confirmed by ELISA at less than 1% cross reactivity against other Ferret heavy or light chain isotypes.

#### **Storage Condition**

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

#### **Precautions Note**



This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Ferret IgM (mu chain) (Peroxidase Conjugated) Secondary Antibody - Protein Information

## Anti-Ferret IgM (mu chain) (Peroxidase Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Ferret IgM (mu chain) (Peroxidase Conjugated) Secondary Antibody - Images