

Mouse IgG3 isotype control Phycoerythrin

Monoclonal MG3 IgG3 , R-Phycoerythrin (RPE) Catalog # ASR3079

Specification

Mouse IgG3 isotype control Phycoerythrin - Product Information

Description

Conjugate Clonality Application Application Note Physical State Host Isotype Buffer

Species of Origin Reconstitution Volume Reconstitution Buffer coniugated **R-Phycoerythrin (RPE)** Monoclonal FC FlowCytometry 1:1000-1:5000 Lyophilized lqG3 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Mouse 1.0 mL Restore with deionized water (or equivalent) 10 mg/mL Bovine Serum Albumin (BSA) -Immunoglobulin and Protease free 0.01% (w/v) Sodium Azide

MOUSE IgG3 isotype control Phycoerythrin

Stabilizer

Preservative

Mouse IgG3 isotype control Phycoerythrin - Additional Information

Shipping Condition Ambient

Purity

This product was prepared from ascitic fluid by affinity chromatography. Assay by immunoelectro-phoresis resulted in a single precipitin arc against anti-Phycoerythrin and Anti-Mouse Serum. Specificity was confirmed by ELISA at less than 1% cross reactivity against other anti-Mouse heavy or light chain isotypes antibodies.

Storage Condition

Store vial at 4° C prior to opening. Dilute only prior to immediate use. Do not freeze after reconstitution. Store reagent in the dark. This product is stable at 4° C as an undiluted liquid. Use subdued lighting during handling and incubation of cells prior to analysis.

Precautions Note This product is for research use only and is not intended for therapeutic or diagnostic applications.

Mouse IgG3 isotype control Phycoerythrin - Protein Information

Mouse IgG3 isotype control Phycoerythrin - Protocols



Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Mouse IgG3 isotype control Phycoerythrin - Images

Mouse IgG3 isotype control Phycoerythrin - Background

Isotype controls are important for Flow Cytometry and have no specificity for target cells within a particular experiment. Their purpose is to confirm the specificity of primary antibody binding that it is not a result of non-specific Fc receptor binding to cells or other cellular protein interactions. Isotype controls need to be matched to the specific primary Abs (species and isotype, including heavy and light chains) being used.