Guinea Pig IgG F(c) Peroxidase
Catalog # ASR3283

Specification

Guinea Pig IgG F(c) Peroxidase - Product Information

Description	**GUINEA PIG IgG F(c) fragment Peroxidase conjugated**
Conjugate	**Peroxidase (Horseradish)**
Physical State	**Lyophilized**
Host Isotype	**IgG F(c)**
Buffer	**0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2**
Species of Origin	**Guinea Pig**
Reconstitution Volume	**1.0 mL**
Reconstitution Buffer	**Restore with deionized water (or equivalent)**
Stabilizer	**10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free**
Preservative	**0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!**

Guinea Pig IgG F(c) Peroxidase - Additional Information

Shipping Condition
Ambient

Purity
This product was prepared from normal serum by delipidation, salt fractionation, ion exchange chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Guinea Pig IgG, anti-Guinea Pig IgG F(c) and anti-Guinea Pig Serum. No reaction was observed against anti-Guinea Pig IgG F(ab')2 or anti-Papain.

Storage Condition
Store vial at 4° C prior to restoration.   For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note
This product is for research use only and is not intended for therapeutic or diagnostic applications.

Guinea Pig IgG F(c) Peroxidase - Protein Information

Guinea Pig IgG F(c) Peroxidase - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Guinea Pig IgG F(c) Peroxidase - Images