

Anti-GM-CSF Secondary Antibody
Rabbit Polyclonal, Unconjugated
Catalog # ASR3291**Specification**

Anti-GM-CSF Secondary Antibody - Product Information

| | |
|------------------|--|
| Description | Anti-GM-CSF (RABBIT) Antibody |
| Host | Rabbit |
| Conjugate | Unconjugated |
| Target Species | Human |
| Reactivity | Human |
| Clonality | Polyclonal |
| Application | ,1,10, |
| Application Note | ELISA 1:1,000-1:5,000;Western Blot 1:500-1:2,000 |
| Physical State | Liquid (sterile filtered) |
| Host Isotype | Antiserum |
| Buffer | None |
| Immunogen | This whole rabbit serum was prepared by repeated immunizations with full length recombinant human GM-CSF. |
| Stabilizer | None |
| Preservative | None |

Anti-GM-CSF Secondary Antibody - Additional Information**Shipping Condition**

Dry Ice

Purity

This antiserum has been heated to 56°C for 30 minutes. In ELISA and other immunoreactive assays, this antiserum will recognize both native and recombinant human GM-CSF in cell supernatants and certain body fluids. This antibody is useful for neutralization of human GM-CSF in bioassays. For neutralization, it is recommended to incubate the sample with a 1:400 dilution of the antiserum for at least 4 hours before being tested. A control of similarly diluted normal rabbit IgG is recommended.

Storage Condition

Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-GM-CSF Secondary Antibody - Protein Information

Anti-GM-CSF Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-GM-CSF Secondary Antibody - Images

Anti-GM-CSF Secondary Antibody - Background

Granulocyte Macrophage Colony Stimulating Factor (also known as GM-CSF, Colony-stimulating factor; CSF, sargramostim and molgramostin) is produced in response to a number of inflammatory mediators by mesenchymal cells present in the hemopoietic environment and at peripheral sites of inflammation. Granulocyte Macrophage-CSF is able to stimulate the production of neutrophilic granulocytes, macrophages, and mixed granulocyte-macrophage colonies from bone marrow cells and can stimulate the formation of eosinophil colonies from fetal liver progenitor cells. GM-CSF can also stimulate some functional activities in mature granulocytes and macrophages. GM-CSF receptors show significant homologies with other receptors for hematopoietic growth factors, including IL2-beta, IL-3, IL-6, IL-7, EPO and the Prolactin receptors.