

Anti-Goat IgG F(c) (Peroxidase Conjugated) Secondary Antibody
Donkey Polyclonal, Peroxidase (Horseradish)
Catalog # ASR3363**Specification****Anti-Goat IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Product Information**

Description	Anti-GOAT IgG F(c) (DONKEY) Antibody Peroxidase Conjugated
Host	Donkey
Conjugate	Peroxidase (Horseradish)
Target Species	Goat
Clonality	Polyclonal
Application	WB, IHC, E
Application Note	ELISA 1:5,000-1:25,000;Western Blot 1:1,000-1:5,000;Immunohistochemistry 1500-1:2,500
Physical State	Lyophilized
Host Isotype	IgG
Target Isotype	IgG F(c)
Buffer	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Immunogen	Goat IgG F(c) fragment
Reconstitution Volume	2.0 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Preservative	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Anti-Goat IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Additional Information**Shipping Condition**

Ambient

Purity

This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Donkey Serum, Goat IgG, Goat IgG F(c) and Goat Serum. No reaction was observed against Goat IgG F(ab')₂.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Goat IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Protein Information

Anti-Goat IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Goat IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Images