

Anti-Human IgG F(c) (Peroxidase Conjugated) Secondary Antibody

Goat Polyclonal, Peroxidase (Horseradish)
Catalog # ASR3385

Specification

Anti-Human IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Product Information

Description Anti-HUMAN IgG F(c) (GOAT) Antibody

Peroxidase Conjugated

Host Goat

Conjugate Peroxidase (Horseradish)

Target Species
Clonality
Application
Human
Polyclonal
WB, E, IC

Application Note ELISA 1:10,000-1:50,000; Western Blot

1:1,000-1:10,000;Immunochemistry

1:500-1:2,500 Lyophilized

Physical State
Host Isotype
IgG
Target Isotype
IgG F(c)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2 Human IgG F(c) fragment

Immunogen Human Reconstitution Volume 2.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Gentamicin Sulfate. Do NOT

add Sodium Azide!

Anti-Human IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Additional Information

Shipping Condition

Ambient

Purity

This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum, Human IgG, Human IgG F(c) and Human Serum. No reaction was observed against Human IgG F(ab')2.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note



This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Human IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Protein Information

Anti-Human IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Human IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Images