

Anti-Dog IgG F(ab')2 Secondary Antibody

Rabbit Polyclonal, Unconjugated Catalog # ASR3510

Specification

Anti-Dog IgG F(ab')2 Secondary Antibody - Product Information

Description

- Host Conjugate Target Species Clonality Application Application Note
- Physical State Host Isotype Target Isotype Buffer

Immunogen Reconstitution Volume Reconstitution Buffer

Stabilizer Preservative Anti-DOG IgG F(ab')2 (RABBIT) Antibody Rabbit Unconjugated Dog Polyclonal ,1,2,10, ELISA 1:20,000-1:100,000;Western Blot 1:2,000-1:10,000;Immunohistochemistry 1:1,000-1:5,000 Lyophilized laG lgG F(ab')2 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Dog IgG F(ab')2 fragment 5.0 mL Restore with deionized water (or equivalent) None 0.01% (w/v) Sodium Azide

Anti-Dog IgG F(ab')2 Secondary Antibody - Additional Information

Shipping Condition Ambient

Purity

This product is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Dog IgG, Dog IgG F(ab')2 and Dog Serum. No reaction was observed against Dog IgG F(c).

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.



Anti-Dog IgG F(ab')2 Secondary Antibody - Protein Information

Anti-Dog IgG F(ab')2 Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Dog IgG F(ab')2 Secondary Antibody - Images